

Award Number: DAMD17-03-1-0402

TITLE: Selective Inhibition of T Cell Tolerance as a Means of  
Enhancing Tumor Vaccines in a Mouse Model of Breast  
Cancer

PRINCIPAL INVESTIGATOR: Jonathan D. Powell, M.D., Ph.D.

CONTRACTING ORGANIZATION: Johns Hopkins University School of  
Medicine  
Baltimore, MD 21205

REPORT DATE: June 2005

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command  
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;  
Distribution Unlimited

The views, opinions and/or findings contained in this report are  
those of the author(s) and should not be construed as an official  
Department of the Army position, policy or decision unless so  
designated by other documentation.

20051101 104

**REPORT DOCUMENTATION PAGE**Form Approved  
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.

**1. REPORT DATE (DD-MM-YYYY)**

01/06/2005

**2. REPORT TYPE**

Annual

**3. DATES COVERED (From -)To)**

15 May 04 - 14 May 05

**4. TITLE AND SUBTITLE**

Selective Inhibition of T Cell Tolerance as a Means of  
Enhancing Tumor Vaccines in a Mouse Model of Breast Cancer

**5a. CONTRACT NUMBER****5b. GRANT NUMBER**

DAMD17-03-1-0402

**5c. PROGRAM ELEMENT NUMBER****6. AUTHOR(S)**

Jonathan D. Powell, M.D., Ph.D.

**5d. PROJECT NUMBER****5e. TASK NUMBER****5f. WORK UNIT NUMBER**

E-Mail: Poweljo@jhmi.edu

**7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)**

Johns Hopkins University School of Medicine  
Baltimore, MD 21205

**8. PERFORMING ORGANIZATION REPORT NUMBER****9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)**

U.S. Army Medical Research and Materiel Command  
Fort Detrick, Maryland 21702-5012

**10. SPONSOR/MONITOR'S ACRONYM(S)****11. SPONSOR/MONITOR'S REPORT NUMBER(S)****12. DISTRIBUTION / AVAILABILITY STATEMENT**

Approved for Public Release; Distribution Unlimited.

**13. SUPPLEMENTARY NOTES****14. ABSTRACT**

Abstract is on page 4.

**15. SUBJECT TERMS**

Breast cancer, vaccine, immunology

**16. SECURITY CLASSIFICATION OF:**

a. REPORT

U

b. ABSTRACT

U

c. THIS PAGE

U

**17. LIMITATION OF ABSTRACT**

UU

**18. NUMBER OF PAGES**

9

**19a. NAME OF RESPONSIBLE PERSON****19b. TELEPHONE NUMBER (include area code)**

## Table of Contents

Cover.....	1
SF 298.....	2
Table of Contents.....	3
Abstract.....	4
Introduction.....	5
Body.....	5
Key Research Accomplishments.....	8
Reportable Outcomes.....	8
Conclusions.....	8
References.....	9

## **Abstract:**

**Purpose:** To determine if the addition of Go6976 to vaccine protocols will inhibit neu specific tolerance and enhance immunotherapy for breast cancer.

**Scope:** In the Her-2/*neu* model of spontaneous breast cancer the immune system of these transgenic mice are tolerant to the neu protein. While immunity to neu can be demonstrated in the neu-transgenic mice (partial breaking of tolerance), this immunity is inadequate in terms of preventing the spontaneous development of tumors and preventing death from tumor challenge. Our experiments involve concomitantly treating mice with the PKC inhibitor Go6976 during tumor vaccine therapy in order to prevent tolerance induction and enhance immunotherapy.

**Findings:** By combining our regimen with a dose of cytoxan we can promote survival of tumor bearing mice when compared with no treatment, vaccine alone or vaccine + cytoxan. We believe that the cytoxan eliminates T regulatory cells. In particular, this combination is very effective in inhibiting tumor growth in the early period post-tumor challenge. Based on these findings we have concludes that extended administration of Go6976 can enhance the efficacy of this regimen even further.

**Significance:** These data support the notion that the novel combination of PKC inhibitor + vaccine can enhance the efficacy of tumor vaccines.

**Introduction:**


In the Her-2/*neu* model of spontaneous breast cancer development it is clear that the immune system of these transgenic mice are tolerant to the neu protein (1-3). In this model not only does the overexpression of neu lead to tumorigenesis but the neu protein is the target of both humoral and cellular immunity which prevent tumor-induced death in the non-transgenic mice (1, 4). Indeed, while immunity to neu can be demonstrated in the neu-transgenic mice (partial breaking of tolerance), this immunity is inadequate in terms of preventing the spontaneous development of tumors and preventing death from tumor challenge. We have demonstrated *in vitro* that the PKC inhibitor Go6976 has the ability to selectively inhibit TCR induced tolerance induction while only minimally inhibiting T cell activation. We hypothesize that the addition of Go6976 to vaccine protocols will inhibit the reinduction of neu specific tolerance and thus facilitate immune mediated protection against the development of spontaneous breast cancer development and tumor challenge.

**Body:****Optimization of drug delivery:**

There was essentially no data concerning *in vivo* use of Go6976. Thus, a major challenge of this proposal (Task 1) was to determine the optimal dose for inhibiting tolerance *neu*-specific tolerance induction in this model. Beginning with an initial educated guess of 5mg/kg every 3 days we have modified the regimen looking for the most effect dose and schedule. As a result of these experiments we have determined that 10mg/kg every 3 days has the ability to enhance vaccine therapy. In addition, it is becoming clear that we get the best results when we continue treatment beyond the initial 3 doses (see Figure 1 below). At this point we are delivering the drug I.P. However, (as mentioned as a possible alternative in our proposal), the data suggest that continuous infusion by ALZET pumps may enhance our effect even more. This remains as a possible future experiment, once we have derived data concerning the optimal delivery I.P.

**Enhancing tumor vaccine therapy by preventing tumor-induced tolerance:**


The basic vaccine protocol is shown in Figure 1 below. Under normal circumstances, tumor vaccines are ineffective in treating tumor in the neu-expressing mice. This is because the T cells from these mice are tolerant to the tumor antigens. The goal of our proposal is to inhibit tumor-induced tolerance pharmacologically with Go6976 and thus enhance the efficacy of the vaccine.

**Figure 1**

Previously, we demonstrated that by inhibiting PKC with Go6976 we could enhance immune function. Therefore a series of experiments were set up to determine if we could enhance vaccine efficacy in terms of survival by adding Go6976 to the vaccine protocol. Interestingly the tumor used for the challenge appeared to be more aggressive. That is, the mice were dying faster than previously reported (data not shown and Figure 3) (5). Of note, Dr. Elizabeth Jaffee's laboratory had been noting the same observations (E. Jaffee personal communication). The enhanced aggressiveness of the tumor did not inhibit our experiments it just meant that the mice were that much more difficult to cure.

In addition, it is becoming increasingly clear that T regulatory cells have the ability to promote tumor-induced tolerance and inhibit vaccine function (6). This has also been found to be true in the *Neu* breast cancer model (Dr. Elizabeth Jaffee personal communication). It has long been known that cytoxan can enhance immune function and it is thought that this is due to the ability of this drug to eliminate T regulatory cells (7). Furthermore, vaccine in the setting of cytoxan have been shown to be marginally effective in this model. Thus, a series of experiments was performed to determine if Go6976 could enhance vaccine therapy + cytoxan.

Mice were treated as described in Figure 1 (10 per group) with either no vaccine, vaccine alone, vaccine+ cytoxan (10mg/kg IP on day 2), or vaccine+cytoxan+Go6976. The mice were evaluated for tumor size and survival.




**Figure 2: Average Tumor size:** Neu-transgenic mice were challenged with tumor on day 0 and then treated as described in the legend above. Cytoxan was given on day 2 while vaccine was given on day 3. Go6976 was given every 3 days (days 6, 9 and 12)

As seen above at day 17 no tumor is evident in the Go6976 treated mice. Of note, there is a statistically significant difference in tumor growth between the vaccine+cytoxan+Go6976 group and the vaccine+cytoxan group. Thus, the effect is not due to the cytoxan but due to the addition of Go6976. The difference is still prevalent on Day 20 but loses significance on Day 22. At this time however, although tumor size in both the vaccine+cytoxan and the vaccine+cytoxan+Go6976 group are equivalent, they are still statistically smaller than the tumors in the vaccine alone group. Thus, as time goes on we lose the "Go6976 effect". Recall, the last dose of Go6976 was on day 12. Thus we interpret these findings as indicating that the addition of Go6976 can enhance vaccine efficacy (Day 17) early and that by continuing Go6976 treatment we might be able to prolong this effect.

In addition to tumor size we also examined the effect of Go6976 on survival. As seen below in Figure 3, the mice treated with vaccine+cytoxan+Go6976 had their survival curve shifted to the right when compared with the other treatments. Furthermore, the curve plateaued at 20% compared with 0% for the other groups. As mentioned above, normally the vaccine treated mice display around 15-30% survival so that for this experiment the conditions were more rigorous. Once again, the marked delay in death leads us to hypothesize that by continuing the treatment we might improve long term survival to a greater extent. We do not believe that the effect of Go6976 is due to the drug itself acting on the tumor because in previous experiments, drug alone did not have any beneficial effect (data not shown).

## Tumor Free survival



**Figure 3: Tumor free survival:** Mice were treated as described for Figure 2, as seen in the figure only the curve for the vaccine+cytoxan+Go6976 treated mice plateaued (20%).

**Key research Accomplishments:**

- \* Proof of the principle that the addition of Go6976 can enhance vaccine efficacy.
- \* Advances in terms of defining optimum dosing.
- \* Observation that a dose of cytoxan + G06976 can synergize in terms of enhancing vaccine therapy

**Reportable outcomes:**

1. Presentation of findings as a poster at "The Era of Hope" meeting

**Conclusions:**

Based on our observations that Go6976 could inhibit anergy induction *in vitro* we hypothesized that this PKC inhibitor could enhance the efficacy of tumor vaccine therapy *in vivo*. Our data clearly show that the addition of Go6976 to the vaccine protocol can delay tumor growth and enhance tumor free survival. These data serve to validate our hypothesis. Interestingly, in our series of experiments the tumor used was more aggressive (perhaps more realistic conditions) than previously published. Thus, these studies were performed under rigorous conditions. Indeed, while vaccine alone normally produces a 15-30% survival rate in our studies it was 0% (5).

An important conclusion from these studies is that we should continue Go6976 therapy for a longer period of time. Indeed, we hypothesize that the marked decrease in tumor size seen on day 17 would be maintained if we continued the drug. As such experiments are planned to test this hypothesis and in doing so further optimize our protocol.

The role of T regulatory cells in promoting tumor-induced tolerance is becoming greater appreciated (6). In this regard the combination of cytoxan and Go6976 appeared to be synergistic. Presumably, the cytoxan helped to eliminate T regulatory cells prior to the vaccine and then Go6976 helped to prevent the reinduction of tolerance after the vaccine. Along these lines, experiments are planned to demonstrate that the decreased tumor size and enhanced survival is due to an increase in tumor-specific T cells. In addition, we will also perform experiments to demonstrate that there is also a decrease in T regulatory cells.

These findings have important implications for the use of vaccines in preventing tumor development (Task 3). Indeed, not surprisingly, the optimal means of overcoming tumor-tolerance will be by inhibiting not only the induction of anergy but also inhibiting T regulatory cells.



## References:

1. Reilly, R. T., M. B. Gottlieb, A. M. Ercolini, J. P. Machiels, C. E. Kane, F. I. Okoye, W. J. Muller, K. H. Dixon, and E. M. Jaffee. 2000. HER-2/neu is a tumor rejection target in tolerized HER-2/neu transgenic mice. *Cancer Res* 60:3569.
2. Rovero, S., A. Amici, E. D. Carlo, R. Bei, P. Nanni, E. Quaglino, P. Porcedda, K. Boggio, A. Smorlesi, P. L. Lollini, L. Landuzzi, M. P. Colombo, M. Giovarelli, P. Musiani, and G. Forni. 2000. DNA vaccination against rat her-2/Neu p185 more effectively inhibits carcinogenesis than transplantable carcinomas in transgenic BALB/c mice. *J Immunol* 165:5133.
3. Kurt, R. A., R. Whitaker, A. Baher, S. Seung, and W. J. Urba. 2000. Spontaneous mammary carcinomas fail to induce an immune response in syngeneic FVBN202 neu transgenic mice. *Int J Cancer* 87:688.
4. Reilly, R. T., J. P. Machiels, L. A. Emens, A. M. Ercolini, F. I. Okoye, R. Y. Lei, D. Weintraub, and E. M. Jaffee. 2001. The collaboration of both humoral and cellular HER-2/neu-targeted immune responses is required for the complete eradication of HER-2/neu-expressing tumors. *Cancer Res* 61:880.
5. Machiels, J. P., R. T. Reilly, L. A. Emens, A. M. Ercolini, R. Y. Lei, D. Weintraub, F. I. Okoye, and E. M. Jaffee. 2001. Cyclophosphamide, doxorubicin, and paclitaxel enhance the antitumor immune response of granulocyte/macrophage-colony stimulating factor-secreting whole-cell vaccines in HER-2/neu tolerized mice. *Cancer Res* 61:3689.
6. Sakaguchi, S., N. Sakaguchi, J. Shimizu, S. Yamazaki, T. Sakihama, M. Itoh, Y. Kuniyasu, T. Nomura, M. Toda, and T. Takahashi. 2001. Immunologic tolerance maintained by CD25+ CD4+ regulatory T cells: their common role in controlling autoimmunity, tumor immunity, and transplantation tolerance. *Immunol Rev* 182:18.
7. Mitchell, M. S. 2003. Combinations of anticancer drugs and immunotherapy. *Cancer Immunol Immunother* 52:686.